

1. Boogart, V. A. 1993-1999. Posidonia Shale Formation ATPO. *DINoloket*.
<https://www.dinoloket.nl/en/posidonia-shale-formation-atpo>.
2. Hess, Hans. 1999. Lower Jurassic Posidonia Shale of Southern Germany. *Fossil Crinoids* **23**:183-196. [http://www.fossilcrinoids.com/literature/9.1_Hess%20\(1999\).pdf](http://www.fossilcrinoids.com/literature/9.1_Hess%20(1999).pdf)
3. Jochum, J., G. Friedrich, D. Leythaeuser, R. Littke, and B. Ropertz. 1995. Hydrocarbon-bearing fluid inclusions in calcite-filled horizontal fractures from mature Posidonia Shale (Hils Syncline, NW Germany). *Ore Geology Reviews* **9**: 363-370
[https://doi.org/10.1016/0169-1368\(94\)00019-K](https://doi.org/10.1016/0169-1368(94)00019-K).
4. Klaver, J., G. Desbois, J. L. Urai, and R. Littke. 2012. BIB-SEM study of the pore space morphology in early mature Posidonia Shale from the Hils area, Germany. *International Journal of Coal Geology* **103**: 12-25. <https://doi.org/10.1016/j.coal.2012.06.012>.
5. Littke, R., D. Leythaeuser, J. Rullkötter and D. R. Baker. 1 January 1991. Keys to the depositional history of the Posidonia Shale (Toarcian) in the Hils Syncline, northern Germany. *Geological Society* **58**: 311-333.
<https://doi.org/10.1144/GSL.SP.1991.058.01.20>
6. Martill, D. 1986. The stratigraphic distribution and preservation of fossil vertebrates in the Oxford Clay of England *Researchgate* **10**: 161-188.
https://www.researchgate.net/profile/David_Martill/publication/291863243_The_stratigraphic_distribution_and_preservation_of_fossil vertebrates_in_the_Oxford_Clay_of_England_UK/links/57835e1708aeb13e7e014774/The-stratigraphic-distribution-and-preservation-of-fossil-vertebrates-in-the-Oxford-Clay-of-England-UK.pdf
7. Montero-Serrano, J., K. B. Föllmi, T. Adatte, J. E. Spangenberg, N. Tribovillard, A. Fantasia, and G. Suan. 2015. Continental weathering and redox conditions during the early Toarcian Oceanic Anoxic Event in the northwestern Tethys: Insight from the Posidonia Shale section in the Swiss Jura Mountains. *Palaeogeography, Palaeoclimatology, Palaeoecology* **429**: 83-99. <https://doi.org/10.1016/j.palaeo.2015.03.043>.
8. Röhl, H., A. Schmid-Röhl, W. Oschmann, A. Frimmel, and L. Schwark. 2001. The Posidonia Shale (Lower Toarcian) of SW-Germany: an oxygen-depleted ecosystem controlled by sea level and palaeoclimate. *Palaeogeography, Palaeoclimatology, Palaeoecology* **165**: 27-52. [https://doi.org/10.1016/S0031-0182\(00\)00152-8](https://doi.org/10.1016/S0031-0182(00)00152-8).

9. Schmid-Röhl, A. and Röhl, H. 2003. Overgrowth on ammonite conchs: environmental implications for the Lower Toarcian Posidonia Shale. *Palaeontology* **46**: 339-352.
[doi:10.1111/1475-4983.00302](https://doi.org/10.1111/1475-4983.00302)
10. Schwark, L. and A. Frimmel. 2004. Chemostratigraphy of the Posidonia Black Shale, SW-Germany: II. Assessment of extent and persistence of photic-zone anoxia using aryl isoprenoid distributions. *Chemical Geology* **206**: 231-248.
<https://doi.org/10.1016/j.chemgeo.2003.12.008>.
11. Song, J., R. Littke, and P. Weniger. 2017. Organic geochemistry of the Lower Toarcian Posidonia Shale in NW Europe. *Organic Geochemistry* **106**: 76-92.
<https://doi.org/10.1016/j.orggeochem.2016.10.014>.